

**WEEKLY TEST TYJ -1 TEST - 25 R**  
**SOLUTION Date 20-10-2019**

**[PHYSICS]**

1. It is the speed of light in free space. Hence, dimension is that of speed, i.e.,  $LT^{-1}$ .

2. Boltzmann's constant = energy/temperature

$$= \frac{[ML^2T^{-2}]}{[\theta]} = [ML^2T^{-2}\theta^{-1}]$$

3. When the body returns to origin, displacement is zero.

$$s = ut + \frac{1}{2}at^2$$

$$0 = 60t - \frac{1}{2} \times 10 \times t^2$$

Solving,  $t = 12$  s

4.  $v \propto \lambda^x \rho^{-y} g^z$

Putting dimensions,

$$LT^{-1} = L^x (ML^{-3})^{-y} (LT^{-2})^z$$

Solving, we get  $v \propto \sqrt{g\lambda}$ .

5. Squaring the given equation,

$$A^2 + B^2 + 2\vec{A} \cdot \vec{B} = C^2$$

Moreover,  $A^2 + B^2 = C^2$  ( $\because A = 6, B = 8, C = 10$ )

$\therefore \vec{A} \cdot \vec{B} = 0$ , i.e.,  $\vec{A}$  is  $\perp$  to  $\vec{B}$

6. Putting equations for  $T$  and  $R$ , we get

$$g \left( \frac{2u \sin \alpha}{g} \right)^2 = 2 \times \frac{u^2 \sin 2\alpha}{g}$$

or  $\tan \alpha = 1$  or  $\alpha = 45^\circ$

7. Let  $t$  be the duration of uniform acceleration. Then,  $(9 - t)$  is the retardation. As the velocity at the end of uniform acceleration and at the beginning of retardation is same, we have

$$0 + at = 0 - 2a \times (9 - t)$$

Solving,  $t = 6$  s

8. The train is moving with horizontal velocity in a straight line, hence vertical ranges will be same.

For a person inside the train, the horizontal range will be zero, because train is an inertial frame. The coin falls back to his hand. For a person outside the train such as  $C$ , the coin has a horizontal velocity and vertical acceleration  $g$ . Hence, it appears to follow a parabolic path. Hence, he observes a horizontal range.



9.

$$R = \frac{u^2 \sin 2\alpha}{g}$$

i.e.,  $R \propto \sin 2\alpha$  (for a given  $u$ )

$$\frac{R_1}{R_2} = \frac{\sin 30^\circ}{\sin 90^\circ} = \frac{1}{2}$$

$$\therefore R_2 = 2R_1 = 4 \text{ km.}$$

10. The acceleration down the plane =  $\frac{g \sin \theta}{1 + (k^2/r^2)}$

This value is maximum when  $k$  is minimum, which happens for a solid sphere ( $k^2 = \frac{2}{5}r^2$ ).

11. Work done = force  $\times$  displacement =  $100 \times \sin 50^\circ \times 1$ .

20% of this work is used to overcome friction. Hence, energy gained = 80% of this work =  $80 \sin 50^\circ$  Joule.

12. The ball moves towards the left due to inertia to a force towards right. This means the bus is taking a right hand turn. The centripetal force is towards the right. Inertial force on the ball (in this case called centrifugal force) acts towards the left.

13. Potential energy of a satellite =  $-\frac{GMm}{R}$

Kinetic energy of a satellite =  $\frac{GMm}{2R}$

$$\therefore \text{Required ratio} = 2 : 1$$

14. According to Kepler's 3rd law,  $T^2 \propto R^3$ .

Now,  $T_1^2 \propto R^3$  (for a satellite close to the earth)

For a geo-synchronous satellite, its distance from the surface of the earth is  $6R$ , and from centre  $7R$ .

So,  $T_2^2 \propto (7R)^3$


$$\therefore T_2 = 7^{3/2} \cdot T_1 = T_1 7\sqrt{7} \text{ or } \frac{T_2}{T_1} = 7\sqrt{7}$$

15. Before the man hits the ground, he is in a state of free fall (ignoring air resistance). A body in a free fall state experiences weightlessness.

16. Here, the disc has only KE of rotation

$$= \frac{1}{2} I\omega^2 = \frac{1}{2} \times \frac{Mr^2}{2} \times \frac{v^2}{r^2} = \frac{1}{4} Mv^2$$

17. Change in velocity = final velocity - initial velocity. It is clear from figure, that change in velocity =  $\sqrt{20^2 + 20^2} = 20\sqrt{2}$  m/s along south-west.



18. KE of a satellite =  $\frac{1}{2} |PE|$

$$KE = \frac{GMm}{2R} \text{ and } PE = -\frac{GMm}{R}$$


$$TE = KE + PE = -\frac{GMm}{2R} = -KE = -2 \times 10^6 \text{ J}$$

19. At the highest point, the projectile has only the horizontal component =  $u \cos \alpha = \frac{u}{2}$  ( $\because \alpha = 60^\circ$ )

$$\text{Range} = \frac{u^2}{g} \sin 2\alpha = \frac{2u^2 \sin \alpha \cos \alpha}{g} = \frac{\sqrt{3}u^2}{2g}$$

20. The cylinder begins to slide when the line through centre of gravity  $G$  passes beyond base of cylinder that is when

$$\tan \theta \geq \frac{r}{h/2} = \frac{2r}{h}$$



Hence,  $h = 2r$  ( $\because \theta = 45^\circ$ )  
 $r = \frac{h}{2} = 5 \text{ cm}$

### **[CHEMISTRY]**

21. (a)  $\text{Na}_2\text{CO}_3 \cdot x\text{H}_2\text{O}$ . Its weight =  $106 + 18x$ .

Weight of anhydrous  $\text{Na}_2\text{CO}_3 = 106$

$$\% \text{ loss in weight} = \frac{18x \times 100}{106 + 18x} = 63$$

$$\therefore x = 10.27 \approx 10$$

22. (c) In law of reciprocal proportions, the two elements combining with the third element, must combine with each other in the same ratio or multiple of that Ratio of S and O when combine with C is 2 : 1. Ratio of S and O is  $\text{SO}_2$ , is 1 : 1

23. (c) Mol in each case

$$7 \text{ g N}_2 = \frac{7}{28} = 0.25; \quad 2 \text{ g H}_2 = \frac{2}{2} = 1.0;$$

$$16 \text{ g NO}_2 = \frac{16}{46} = 0.34; \quad 16 \text{ g O}_2 = \frac{16}{32} = 0.50$$

Thus hydrogen has maximum moles, hence maximum molecules.

- 24.

$$(a) \lambda = \frac{h}{mv} = \frac{h}{\sqrt{2mE}}$$


$$= \frac{6.6 \times 10^{-34}}{\sqrt{2 \times 1 \times 0.5}} = 6.6 \times 10^{-34}$$




25. (b) 1.  $\left. \begin{array}{l} \text{BO}_3^{3-} \longrightarrow 5 + 8 \times 3 + 3 = 32 \\ \text{CO}_3^- \longrightarrow 6 + 8 \times 3 + 2 = 32 \\ \text{NO}_3^- \longrightarrow 7 + 8 \times 3 + 1 = 32 \end{array} \right\} \text{ISO electronic}$
2.  $\left. \begin{array}{l} \text{SO}_3^- \longrightarrow 16 + 8 \times 3 + 2 = 42 \\ \text{CO}_3^- \longrightarrow 32 \\ \text{NO}_3^- \longrightarrow 32 \end{array} \right\} \text{not ISO electronic}$
3.  $\left. \begin{array}{l} \text{CN}^- \longrightarrow 6 + 7 + 1 = 14 \\ \text{N}_2 \longrightarrow 7 \times 2 = 14 \\ \text{C}_2^- \longrightarrow 6 \times 2 + 2 = 14 \end{array} \right\} \text{ISO electronic}$
4.  $\left. \begin{array}{l} \text{PO}_4^{3-} \longrightarrow 15 \times 8 + 4 + 3 = 50 \\ \text{SO}_4^- \longrightarrow 16 + 8 + 2 = 50 \\ \text{ClO}_4^- \longrightarrow 17 + 8 \times 4 + 1 = 50 \end{array} \right\} \text{ISO electronic}$

26. (c)  $ns^2 p^1$  is the electronic configuration of III period.  
 $\text{Al}_2\text{O}_3$  is amphoteric oxide

27. (c)  $\text{PF}_5$  trigonal bipyramidal



$\text{BrF}_3$  square pyramidal (distorted)





28. (d)  $\begin{array}{c} (-) \\ \text{:}\ddot{\text{O}}\text{:} - \text{N} \begin{array}{l} \nearrow \ddot{\text{O}}\text{:} \\ \searrow \ddot{\text{O}}\text{:} \end{array} \end{array}$  It has 4 bond pairs and none lone pair on N.

29. (d) The value of  $a$  is a measure of the magnitude of the attractive forces between the molecules of the gas. Greater the value of 'a', larger is the attractive intermolecular force between the gas molecules. The value of  $b$  related to the effective size of the gas molecules. It is also termed as excluded volume. The gases with higher value of  $a$  and lower value of  $b$  are more liquefiable, hence for  $\text{Cl}_2$  "a" should be greater than for  $\text{C}_2\text{H}_6$ , but for it  $b$  should be less than for  $\text{C}_2\text{H}_6$ .

30. (d)  $\text{NH}_3$  and  $\text{HCl}$  react to form  $\text{NH}_4\text{Cl}$
31. (b) This is combustion reaction, which is always exothermic hence  
 $\Delta H = -ve$   
 As the no. of gaseous molecules are increasing hence entropy increases  
 Now  $\Delta G = \Delta H - T\Delta S$   
 For a spontaneous reaction  
 $\Delta G = -ve$   
 Which is possible in this case as  $\Delta H = -ve$  and  $\Delta S = +ve$ .

32. (c)  $\sigma 1s^2, \sigma^* 1s^2, \sigma 2s^2, \sigma^* 2s^2,$   
 $\sigma 2p_z^2, \pi 2p_x^2, \pi 2p_y^2, \pi^* 2p_x^2, \pi^* 2p_y^2$   
 $\therefore$  No. of antibonding electron pairs = 4

33. (d) The shape of  $\text{BF}_3$  is trigonal planar  and  $\mu = 0$  hence it is non polar. The shape of  $\text{NF}_3$  is  pyramidal  $\delta^- \text{F} \text{N}^{\delta+} \text{F} \delta^-$  and  $\mu \neq 0$  hence it is polar.

34. 10% (w/w) solution means  $100 \text{ g} = \frac{100}{1.1} \text{ ml}$   
 solution contains 10 g solute,  $\text{NaOH}$   

$$\text{Molarity} = \frac{w \times 1000}{m^l \times v} = \frac{10 \times 1000}{40 \times \left(\frac{100}{1.1}\right)} = 2.75 \text{ M}$$

35. B
36. [a] The average energy per bond in  $\text{O}_2$  is greater than that in  $\text{O}_3$  because dissociation of  $\text{O}_2$  is endothermic